



## Orbital Tori Construction Using Trajectory Following Spectral Methods

---

By Ralph E. Bordner

Biblioscholar Okt 2012, 2012. Taschenbuch. Book Condition: Neu. 246x189x11 mm. This item is printed on demand - Print on Demand Neuware - The Kolmogorov-Arnold-Moser (KAM) theorem states that lightly perturbed integrable Hamiltonian systems maintain their multiply periodic, toroidal motion in the phase space. The assertion that earth orbiting satellites under the influence of the geopotential mimic this behavior is the underlying premise of this work. This paper focuses on applying trajectory following spectral methods on selected orbits to decompose them into multi-periodic Fourier series, effectively compressing ephemerides for long-term use. The proposed approach focuses on fitting local spectral structures, denoted as frequency clusters, within the sampled orbital data to the analytical form of the windowed, truncated, continuous Fourier transform. This approach is significantly more numerically efficient than fitting every coefficient within the N-tuple Fourier series simultaneously. Numerical results using integrated data show maximum error in orbital torus fits can be kept at the single-digit-meter level per coordinate axis over a 1 year period. 188 pp. Englisch.



**READ ONLINE**  
[ 7.32 MB ]

### Reviews

*This publication is amazing. It is definitely basic but shocks in the fifty percent of your publication. You wont feel monotony at anytime of your own time (that's what catalogues are for concerning if you question me).*

*-- Prof. Kirk Cruickshank DDS*

*This kind of book is every little thing and taught me to looking ahead of time and a lot more. I am quite late in start reading this one, but better then never. I found out this book from my dad and i encouraged this pdf to find out.*

*-- Justus Hettinger*